

[DISPENSER PACK LABEL – 30 g, 60 g TUBES]

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

			
Maxforce[®]			
COCKROACH GEL			
ACTIVE CONSTITUENT: 20 g/kg HYDRAMETHYLNON			
	GROUP	20	INSECTICIDE

*** g
Net**

A gel bait for the control of cockroaches by professional pest control operators as per the directions for use.

Before using this product read all directions on outer pack. This dispenser must be returned to the outer pack after use. Do not destroy outer pack while product still remains.
 Sale of this dispenser except in unopened pack is illegal.

(label code)

***30 g, 60 g**

[DISPENSER PACK LABEL – 30 g, 60 g TUBES] CONTD

DIRECTIONS FOR USE:

PEST: For the control of German and other common cockroaches.

RATE: Apply as 1-2 spots or in a band 75 to 150 mm long per square metre.

Recommended spot size $\frac{1}{4}$ to $\frac{1}{2}$ g.

NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

STORAGE AND DISPOSAL:

Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight. Dispose of empty container by wrapping in paper, placing in plastic bag and putting in the garbage.

SAFETY DIRECTIONS:

Harmful if swallowed. When using this product wear rubber gloves. Wash hands after use. After each day's use wash gloves.

FIRST AID:

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone Australia: 13 11 26).

Batch Number:

Date of Manufacture:

APVMA Approval Number: 56626/53378

Bayer Environmental Science
A Business Group of Bayer CropScience Pty Ltd
A.B.N. 87 000 226 022
391-393 Tooronga Rd
East Hawthorn, Vic. 3123
Phone: (03) 9248 6888
Fax: (03) 9248 6800
Website: www.bayeres.com.au

(OUTER PACK LABEL – MAIN PANEL 30 g, 60 g, 4x30 g)

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

			
Maxforce[®]			
COCKROACH GEL			
ACTIVE CONSTITUENT: 20 g/kg HYDRAMETHYLNON			
	GROUP	20	INSECTICIDE

*** g Net**

A gel bait for the control of cockroaches by professional pest control operators as per the directions for use.

30 and 60 g: Contains one measure pack which it is illegal to sell separately.

4x30 g: This pack contains 4 pre-filled reservoirs for use with Maxforce Bait Injector. Sale of these dispensers except in unopened pack is illegal.

(label code)

* 30 g, 60 g, 4x30 g

(OUTER PACK - ANCILLARY PANEL LABEL 30 g, 60 g, 4x30 g)**DIRECTIONS FOR USE**

Situations	Pest	Rate
Barracks, Farm buildings, Food processing establishments, Factories, Homes, Hospitals, Meatworks (non- product areas), Offices, Restaurants, Schools, Shops, Storerooms, Transport vehicles (including aircrafts) & facilities.	For the control of German and other common cockroaches.	Apply as 1-2 spots or in a band 75 to 150 mm long per square metre. Recommended spot size ¼ to ½ g. DO NOT place in areas that are routinely washed.

NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

GENERAL INSTRUCTIONS:**30 g only: DIRECTION FOR LOADING RESERVOIRS and OPERATION:**

On the handle of the drive rod point the arrow facing up. Pull the drive rod as far as it will go. Attach a pre-filled reservoir of MAXFORCE gel to the bait injector with a twist locking movement. Break off and dispose of the plastic tip and attach the dispensing cap. Push the drive rod until the plunger seats against the piston in the reservoir of the gel. Turn the drive rod knob until teeth are facing down and engaging the trigger mechanism (arrow down). Gently squeeze the trigger until the gel flows out.

30 g only: DIRECTIONS FOR CHANGING RESERVOIRS:

With the arrow on the drive rod pointing up, pull the drive rod out as far as it will go. Remove the empty reservoir, wrap and place in rubbish bin. Reservoirs that still contain gel can be sealed with the supplied cap for further use. To attach a new reservoir, follow the directions above.

60 g only: To apply gel, remove the cap on the nozzle, touch the tip to a surface and depress the plunger. Recap the dispenser after treatment is completed.

Maxforce Cockroach Gel is an insect stomach poison. The bait will adhere to non-greasy and non-dusty surfaces and will remain pliable and palatable to cockroaches as long as it is visibly present. Re-apply when bait is no longer visibly present, according to the remaining level of infestation. A visible inspection of bait placements is recommended at one month after the initial treatment. Re-application should occur if the bait is visibly depleted. Pest population reduction will be apparent within a week or less. Avoid contact with porous surfaces as the bait may stain.

INSECTICIDE RESISTANCE WARNING

GROUP	20	INSECTICIDE
--------------	-----------	--------------------

For insecticide resistance management Maxforce Cockroach Gel is a Group 20 insecticide. Some naturally occurring insect biotypes resistant to Maxforce Cockroach Gel and other Group 20 insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if Maxforce Cockroach Gel or other Group 20 is used repeatedly. The effectiveness of Maxforce Cockroach Gel on resistant individuals could be significantly reduced. Since the occurrence of resistant

individuals is difficult to detect prior to use, Bayer CropScience Pty Ltd accepts no liability for any losses that may result from failure of Maxforce Cockroach Gel to control resistant insects. For further information on managing insecticide resistance contact your local supplier or Bayer Environmental Science representative.

PRECAUTIONS:

Avoid contact with food, food utensils or places where food is prepared or stored. Do not apply on food preparation or serving areas. Should not be used in areas accessible to children. Do not use on or around electrical equipment due to the possibility of electric shock. DO NOT apply the product in areas easily accessible to children and animals.

STORAGE AND DISPOSAL:

Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight. Dispose of used tube by wrapping in paper, placing in plastic bag and putting in garbage.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT:

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

SAFETY DIRECTIONS:

Harmful if swallowed. When using the product wear rubber gloves. Wash hands after use. After each day's use wash gloves.

FIRST AID:

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone Australia: 13 11 26).

MATERIAL SAFETY DATA SHEET

Additional information is listed in the material safety data sheet.

Exclusion of Liability

This product must be used strictly as directed, and in accordance with all instructions appearing on the label and in other reference material. So far as it is lawfully able to do so, Bayer CropScience Pty Ltd accepts no liability or responsibility for loss or damage arising from failure to follow such directions and instructions.

Maxforce® is a registered trademark of Bayer.
APVMA Approval Number: 56626/53378

Bayer Environmental Science
Bayer CropScience Pty Ltd
A.B.N. 87 000 226 022
391-393 Tooronga Rd
East Hawthorn, Vic. 3123
Fax: (03) 9248 6800

Phone: (03) 9248 6888
Website: www.bayeres.com.au

BATCH NO:

DATE OF MANUFACTURE:

BAR CODE