

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

MAXFORCE® ACTIV COCKROACH GEL

Active Constituent: 10 g/kg CLOTHIANIDIN

GROUP	4A	INSECTICIDE
--------------	-----------	--------------------

For the control of cockroaches in domestic, commercial, public and industrial situations as specified in the Directions for Use.

Before using product, read directions on the outer pack. Sale of this syringe except in outer pack is illegal.

Net contents: 30 g

Bayer CropScience Pty Ltd
Level 1, 8 Redfern Rd,
Hawthorn East, Vic. 3123
Telephone: (03) 9248 6888
Technical enquiries: 1800 804 479
www.environmentalscience.bayer.com.au

STORAGE AND DISPOSAL Store in the closed, original container in a cool dry place. DO NOT store in direct sunlight. Completely empty syringe in normal use. Dispose of empty syringe and packaging by wrapping in paper, placing in plastic bag and putting in garbage.

SAFETY DIRECTIONS Wash hands after use.

FIRST AID If poisoning occurs, contact a doctor or Poisons Information Centre (Phone 13 11 26).

APVMA Approval No.: 81040 / 102203

Batch number

Date of Manufacture

HAACP logo to be confirmed

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

MAXFORCE® ACTIV COCKROACH GEL

Active Constituent: 10 g/kg CLOTHIANIDIN

GROUP	4A	INSECTICIDE
--------------	-----------	--------------------

For the control of cockroaches in domestic, commercial, public and industrial situations as specified in the Directions for Use.

The syringe must not be sold separately. Do not destroy outer pack while product still remains.

Net contents: 90 g (3 x 30 g)

Bayer CropScience Pty Ltd
Level 1, 8 Redfern Rd,
Hawthorn East, Vic. 3123
Telephone: (03) 9248 6888
Technical enquiries: 1800 804 479
www.environmentalscience.bayer.com.au

STORAGE AND DISPOSAL Store in the closed, original container in a cool dry place. DO NOT store in direct sunlight. Completely empty syringe in normal use. Dispose of empty syringe and packaging by wrapping in paper, placing in plastic bag and putting in garbage.

SAFETY DIRECTIONS Wash hands after use.

FIRST AID If poisoning occurs, contact a doctor or Poisons Information Centre (Phone 13 11 26).

SAFETY DATA SHEET Additional information is provided in the Safety Data Sheet which may be found at www.environmentalscience.bayer.com.au

APVMA Approval No.: 81040 / 102203

DIRECTIONS FOR USE

SITUATIONS	PEST	RATE	CRITICAL COMMENTS
<p>Domestic premises such as houses and residential buildings including kitchens;</p> <p>Food manufacture and processing establishments, kitchens and abattoirs;</p> <p>Food storage areas such as in food retailing establishments, warehouses, raw material stores, silos except where grain is stored;</p> <p>Food preparation areas such as in restaurants and commercial kitchens;</p> <p>Public buildings including hospitals, nursing homes, schools, hotels, public baths, municipal buildings, churches, halls, community centres, airports, train or bus stations, cinemas, shops etc.;</p> <p>Small scale animal housing such as zoos, pet shops, kennels, veterinary practices, laboratory animal houses etc. (excluding animal pens and cages);</p> <p>Modes of transportation such as buses, trucks, trailers, rail cars and marine vessels.</p>	<p>Small cockroaches e.g. German cockroach (<i>Blatella germanica</i>); Brown-banded cockroach (<i>Supella longipalpa</i>)</p>	<p><u>Light to moderate infestations or retreatment:</u> One placement per m²;</p> <p><u>Heavy infestations:</u> Two placements per m²</p>	<p>One placement of 0.1 g is equivalent to a 7 mm diameter spot. A thin band equating to one placement of 0.1 g may alternatively be applied. The application rate is dependent on the level of infestation and species to be controlled. A moderate infestation is when cockroaches are rarely visible during the day. A heavy infestation is when cockroaches are commonly visible during the day.</p> <p>Apply Maxforce Activ directly as spots or thin bands to surfaces in identified target areas. The product should be placed out of sight and where light intensity is low. Place the bait placements out of reach of children or pets. Avoid application to excessively dusty, damp or greasy locations and areas which are subject to washing. Do not apply to areas that have recently been treated with other insecticides or contaminate the bait with other insecticides.</p> <p>Reassessment of treated areas one month after initial baiting is recommended to evaluate if additional baiting is required. Bait placements should be replenished when consumed or at least every three months as long as pest populations persist.</p>
	<p>Large cockroaches e.g. American cockroach, Australian cockroach (<i>Periplaneta</i> spp.); Oriental cockroach (<i>Blatta orientalis</i>)</p>	<p><u>Light to moderate infestations or retreatment:</u> Two placements per m²;</p> <p><u>Heavy infestations:</u> Three placements per m²</p>	

NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

DO NOT allow to contact food, livestock feed or their water. DO NOT place gel on surfaces where food is prepared or on utensils that are used for food preparation, on food packing materials or on surfaces where food packing materials are stored. DO NOT place gel on surfaces which are regularly washed. Gel should be applied out of reach of children, pets and other animals.

GENERAL INSTRUCTIONS

APPLICATION Apply this product using a suitable application device. To apply the gel, remove the cap on the nozzle and touch the tip to the surface to be treated. Slowly depress plunger until sufficient gel is dispensed. When the treatment has been completed remove the empty syringe and dispose of safely. If the syringe is only partially used, remove it from the gun, seal it with the end cap provided and store as directed. Unwanted gel may be removed by sponging with a 5% sodium chloride (table salt) solution.

RESISTANCE MANAGEMENT For insecticide resistance management, Maxforce Activ Cockroach Gel is a Group 4A insecticide. Some naturally occurring insect biotypes resistant to Maxforce Activ and other Group 4A insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if Maxforce Activ or other Group 4A insecticides are used repeatedly. The effectiveness of Maxforce Activ on resistant individuals could be significantly reduced. Since occurrence of resistant individuals is difficult to detect prior to use, Bayer CropScience Pty Ltd. accepts no liability for any losses that may result from the failure of Maxforce Activ to control resistant insects. Maxforce Activ may be subject to specific resistance management strategies. For further information contact your local supplier or Bayer Environmental Science representative.

PRECAUTIONS Warning: Some sensitive materials e.g. textiles may be stained by the product. If in doubt, apply product to a small, hidden area first. DO NOT attempt to open or refill syringe.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT DO NOT contaminate streams, rivers or waterways with the gel or used syringes.

STORAGE AND DISPOSAL: Store in the closed, original container in a cool dry place. DO NOT store in direct sunlight. Completely empty syringe in normal use. Dispose of empty syringe and packaging by wrapping in paper, placing in plastic bag and putting in garbage.

SAFETY DIRECTIONS: Wash hands after use.

FIRST AID: If poisoning occurs, contact a doctor or Poisons Information Centre (Phone Australia 13 11 26).

APVMA Approval No.: 81040 / 102203

Exclusion of Liability: This product must be used strictly as directed, and in accordance with all instructions appearing on the label and in other reference material. So far as it is lawfully able to do so, Bayer CropScience Pty Ltd accepts no liability or responsibility for loss or damage arising from failure to follow such directions and instructions.

Bayer CropScience Pty Ltd
Level 1, 8 Redfern Rd,
Hawthorn East, Vic. 3123
Telephone: (03) 9248 6888
Technical enquiries: 1800 804 479
www.environmentalscience.bayer.com.au

©Maxforce is a registered trademark of the Bayer Group.

MAXFORCE ACTIV COCKROACH GEL

81040/102203

LABEL FOR OUTER PACK

Batch Number

Date of Manufacture

HAACP logo to be confirmed

GHS Classification not required on this label